

# Are health inequalities increasing in Japan? The trends of 1955 to 2000

Yoshiharu Fukuda\*, Hiroyuki Nakao, Yuichiro Yahata, Hirohisa Imai

Department of Epidemiology, National Institute of Public Health, 2-3-6 Minami, Wako, Saitama, Japan.

**SUMMARY** This study aimed to elucidate trends in socioeconomic inequalities in health during the past half century in Japan. Association of life expectancy and age-adjusted mortality with per capita income was examined using data on prefectures and municipalities in Japan of 1955 to 2000 via the slope index of inequality (SII) and Poisson regression. Although there were a few differences among health indicators and sex, health inequalities by prefecture, measured by the SII, decreased from 1955 to 1995. However, health inequalities increased from 1995 to 2000 both for life expectancy and mortality. Similar trends were found in municipal analyses: the association between income and mortality, measured by the rate ratio from Poisson regression, decreased until 1995 but increased from 1995 to 2000. In the past half century, and especially until 1995, geographical health inequalities decreased in Japan, while from 1995 to 2000 health inequalities appeared to increase. Recent social conditions including the possible increase in social inequalities may have contributed to this increase. Careful monitoring and elimination of social and health inequalities should be encouraged.

**Key Words:** Health inequalities, socioeconomic factors, life expectancy, ecological study

## Introduction

Elimination of health inequalities has been a great challenge in international and domestic public health policy. A large number of studies have demonstrated health inequalities attributable to socioeconomic conditions, including income, educational attainment, social class, and other factors (1-4). The degree of socioeconomic inequalities in society is closely linked to the health of the population (5,6).

Japan has shown marked improvement in the health of the population in the past half century. Major health indicators such as life expectancy and infant mortality have been ranked as some of the world's highest (7). In addition to economic growth and improved living standards, decreased socioeconomic inequalities and an egalitarian social system are considered to contribute to the health improvement of Japanese (6,8-11).

This egalitarian society, however, may be changing. Researchers in the fields of economics, sociology, and

education are extremely concerned about increasing socioeconomic inequalities in Japan, and especially in the past decade (12-14). Although more discussion is needed, the social conditions underlying the increasing inequalities include economic recession and recent economic, taxation, and social security policies (12-14). Little is known about the influence of the possible increase in socioeconomic inequalities in health, leading to the question of if health inequalities are increasing in Japan.

This study elucidated the trends in health inequalities during the past half century in Japan. To this end, an ecological approach was taken at the prefectural and municipal levels to gather data in order to facilitate further debate on health inequalities.

## Methods

### *Populations studied and observation period*

The populations studied were prefectures and municipalities. These are basic administrative divisions in Japan: the prefecture is the higher level and consists of municipalities. There are currently 47 prefectures, an increase from 46 after 1972 with the reversion of

\*Correspondence to: Department of Epidemiology, National Institute of Public Health, 2-3-6 Minami, Wako, Saitama 351-0197, Japan; e-mail: fukuday@niph.go.jp

Received May 9, 2007  
Accepted May 24, 2007

**Table 1.** Mean and coefficient of variance (CV) of per capita income, life expectancy, and age-adjusted mortality of 46 prefectures in Japan, 1955 to 2000

Year	Per capita income (thousand yen)		Life expectancy (years)				Age-adjusted mortality (per 100,000)			
	Mean	(CV)	Men		Women		Men		Women	
			Mean	(CV)	Mean	(CV)	Mean	(CV)	Mean	(CV)
1955	68.7	(22.0)	62.95	(2.03)	67.13	(1.93)	-		-	
1960	112.2	(26.6)	65.19	(1.66)	70.07	(1.37)	-		-	
1965	215.5	(23.4)	67.49	(1.44)	72.88	(1.01)	632.0	(9.31)	375.9	(6.37)
1970	495.4	(21.7)	69.41	(1.35)	75.05	(0.71)	566.7	(10.64)	321.8	(5.84)
1975	1019.1	(14.2)	71.44	(1.06)	76.84	(0.64)	459.1	(9.65)	255.0	(5.50)
1980	1575.0	(12.7)	73.34	(0.92)	78.92	(0.54)	402.0	(9.92)	206.2	(6.06)
1985	1960.9	(14.6)	74.82	(0.80)	80.73	(0.53)	365.2	(9.50)	176.6	(6.46)
1990	2634.7	(15.9)	75.96	(0.78)	82.10	(0.47)	335.5	(9.01)	155.2	(5.57)
1995	2865.5	(13.2)	76.66	(0.76)	83.30	(0.54)	319.9	(10.38)	148.9	(9.12)
2000	2867.6	(12.4)	77.62	(0.75)	84.70	(0.48)	297.7	(8.82)	135.6	(6.28)

CV = Standard deviation/Mean × 100

Okinawa Prefecture to Japan. Okinawa Prefecture was excluded from the current analyses to ensure comparability of time trends.

Municipalities include cities (“shi”), towns (“machi”), villages (“mura”), and wards (“ku”). The number of municipalities fluctuated and numbered almost 3350 during the observed periods because of mergers and dissolutions of municipalities.

The entire observation period was 1955 to 2000, but the analytical period depended upon variables because of limited data availability.

*Data*

Health indicators were life expectancy (LE) and mortality. In prefectural analyses, LE and age-adjusted mortality among populations aged 20 to 64 years were used. The data were obtained from the Prefectural Life Table and Vital Statistics (15-18).

In municipal analyses, the observed number of deaths was obtained from Vital Statistics (19,20) and aggregated in intervals of five consecutive years (1973-77, 1978-82, 1983-87, 1988-92, 1993-97, and 1998-2000). The expected number of deaths was estimated using the age-specific population of the municipality and age-specific mortality of the entire country (19-21). Analysed municipalities numbered 3346, 3348, 3356, 3346, 3361, and 3356, respectively, for the five observation periods.

Per capita income served as a socioeconomic indicator. Per capita income by prefecture and municipality was obtained from a published database (21,22).

*Analyses*

In prefectural analyses, the slope index of inequality (SII) served as a measure of the association between health indicators and income. The SII is estimated from the slope of the linear regression line between income

ranking and health indicator and the mean of the health indicator (23). Since the SII is independent of absolute values of health and its predictive variables, it is useful for comparison of the magnitude of health inequalities, and especially for comparing time trends and different indicators (23).


First, the prefectures were ranked according to per capita income, and ranking values ranging from 0 (lowest income) to 1 (highest income) were assigned to prefectures. Then, the linear regression line was fitted with the health indicator as the dependent variable and the ranking value as the explanatory variable. The SII was estimated by dividing the slope by the mean of the health indicators and presented as a percentage (× 100). The observation period was 1955 to 2000 for LE and 1965 to 2000 for mortality.

In municipal analyses, Poisson regression was used with the number of observed and expected deaths and per capita income, and the rate ratio (RR) of income for mortality was estimated. Two income variables were separately introduced. First, per capita income was introduced as a continuous variable (in units of a million yen). Second, an ordinal variable was used: the lowest decile = 0.05 to the highest decile = 0.95. The observation period was 1975 to 2000. Municipal per capita income in 1975 was not available, so that in 1980 was used for the 1975 analysis.

SPSS 15.0J was used for linear regression analysis and MLwiN 2.02 for Poisson regression analysis.

**Results**

Table 1 summarizes the health indicators and per capita income of 46 prefectures from 1955 to 2000. Income markedly increased, and especially until 1990, with narrowing of the variation in accordance with the coefficient of variance (CV). LE continuously increased with narrowing variation. Age-adjusted mortality continuously decreased but was not accompanied by narrowing variation.




**Figure 1.** Slope index of inequality for income and life expectancy for 46 prefectures in Japan, 1955 to 2000. Vertical lines show the 95% confidence interval.

Figure 1 shows the SII for LE and per capita income in prefectural analyses from 1955 to 2000. A positive sign means that a prefecture with higher incomes had a longer LE. The SII decreased during 1955 to 1995 for women and during 1970 to 1995 for men. In contrast, the SII increased for both men and women from 1995 to 2000.

The SII for age-adjusted mortality and per capita income from 1965 to 2000 is shown in Figure 2. A negative sign means that a prefecture with higher incomes had a lower mortality rate. According to the SII, the association between income and mortality decreased until 1995 but increased from 1995 to 2000.

Table 2 shows the results of municipal analyses, demonstrating the RR of per capita income for mortality. An RR of less than 1.0 means that municipalities with higher incomes had a lower mortality rate. There were some differences in absolute values between continuous and categorical variables due to the different units, but the time trends were similar. The association between income and mortality decreased until 1995. For women in 1990 and 1995, the RR was more than 1.0, showing that municipalities with higher incomes had a higher mortality rate. From 1995 to 2000, RR increased in both



**Figure 2.** Slope index of inequality for income and age-adjusted mortality for 46 prefectures in Japan, 1965 to 2000. Vertical lines show the 95% confidence interval.

men and women.

### Discussion

This paper demonstrated the possibility of health inequalities increasing from 1995 to 2000 in Japan. An increase in health inequalities over the past few decades has been found in other industrial countries (24-26). The current findings offer the first suggestion of a recent increase in health inequalities in Japan.

Until 1995, the association between income and health indicators decreased, as shown in previous studies (8,9,27). The decrease in health inequalities was accompanied by significant LE extension and a decline in mortality. Possible contributors to the improved health of the Japanese population have been noted. An egalitarian social system and culture appears to contribute substantially through compulsory education, universal health insurance coverage, public health services, income adjustment policy, and strong social relationships (6,8-10). This is considered to be a good indication that fewer socioeconomic inequalities improve the health of the population (6,11).

The association of income and health indicators

**Table 2.** Results of poisson regression of per capita income and mortality by municipal level: rate ratio with 95% confidence interval

Year	Men		Women	
	Continuous <sup>a</sup>	Categorical <sup>b</sup>	Continuous <sup>a</sup>	Categorical <sup>b</sup>
1975	0.779 (0.766, 0.793)	0.861 (0.851, 0.872)	0.877 (0.861, 0.893)	0.945 (0.933, 0.958)
1980	0.855 (0.841, 0.870)	0.920 (0.909, 0.931)	0.938 (0.921, 0.955)	0.987 (0.975, 1.000)
1985	0.876 (0.865, 0.887)	0.913 (0.902, 0.924)	0.959 (0.946, 0.972)	0.998 (0.878, 1.134)
1990	0.950 (0.943, 0.957)	0.941 (0.940, 0.942)	1.004 (0.996, 1.012)	1.042 (1.030, 1.055)
1995	0.944 (0.936, 0.951)	0.935 (0.925, 0.945)	1.012 (1.003, 1.020)	1.035 (1.023, 1.048)
2000	0.886 (0.883, 0.889)	0.853 (0.849, 0.857)	0.975 (0.972, 0.979)	0.972 (0.967, 0.977)

<sup>a</sup>Per capita income was used as the continuous variable in units of a million yen.

<sup>b</sup>Per capita income was used as the ordinal variable: from the lowest decile of 0.05 to the highest decile of 0.95.

increased from 1995 to 2000. This increase was consistent regardless of different health indicators and different geographic levels for both men and women. The recent data invite several warnings about Japanese health status. LE of some occupational classes declined in the past few years (28). LE of all Japanese men also declined from 2004 to 2005 (29). This is not conclusive, but the increase in health inequalities may be linked to the deterioration of the health of the population.

Although the explanation for the possible increase in health inequalities in recent years is beyond the scope of this study, increasing socioeconomic inequalities are a potential contributor to increasing health inequalities. Some measures such as the Gini coefficient suggest a widening of income distribution in Japan (14,30). The economic recession after the collapse of the bubble economy in the early 1990s and the subsequent policies on economics, taxation, and social security might have contributed to increased socioeconomic inequalities (12-14). Crumbling of the lifetime employment system found in Japanese companies, the increase in unstable employment, and the increase in social security costs might have also accelerated worries about increasing socioeconomic inequalities (12-14).

The health care system in Japan is considered among the best in the world in terms of fairness of financial contribution, health outcomes, and other indicators (31). The system is believed to contribute to the healthy status of the Japanese population (9). However, recent figures suggest an increase in inequality in access to and use of health care in Japan. Geographic disproportions in health care, such as in the number of obstetricians, gynecologists and pediatricians, and cancer care resources are increasingly receiving attention (32,33). A previous ecological study showed that the lack of resources for maternal and child health is associated with higher infant mortality in Japan (34). Other studies have noted that the postgraduate medical training system and recent health policies, mainly in relation to the postgraduate medical training system and control of health care expenditures, might trigger geographic disproportions and widening inequalities in health care (35). In addition, an increasing number of people who cannot afford insurance premiums appears to be endangering universal health insurance coverage (36). Circumstances concerning health care may widen health inequalities and consequently threaten improvements in the health of the Japanese population.

Analyses of health inequalities often suffer from methodological problems. The selection of both health indicators and socioeconomic variables and methods of analyzing their association are critical (37). The findings of this study were obtained using sophisticated methods with reliable health and socioeconomic variables at two different levels. Nonetheless, a few limitations are acknowledged below.

First, the observation period is too short to conclude

that health inequalities were increasing until 1995, and health inequalities should be continuously monitored. Second, another combination of health indicators and socioeconomic variables could demonstrate a different pattern from that of this study. More specified health indicators, such as cause-specific mortality, will elucidate more detailed situations including an explanation for increasing health inequalities. Area indicators representing socioeconomic conditions are critical in area-based analyses. Agreed-on area indicators have not been established in Japan, unlike in some countries where indicators such as deprivation indices have been applied (38). The development of area-based socioeconomic indicators is an urgent challenge for the study of health inequalities in Japan. Lastly, but of equal importance, ecological studies have methodological limitations, including confounding factors and the ecological fallacy (39). Nonetheless, geographical data can yield meaningful evidence on health inequalities, especially in the long term, since individual-level data are generally of limited use for such analyses (40). Further studies with individual-level analyses based on a system to monitor individual-level inequalities should be encouraged in order to provide more conclusive evidence.

In conclusion, this study showed a possible increase in socioeconomic inequalities in life expectancy and mortality from 1955 to 2000, following a decrease in inequalities from 1955 to 1995. Although conclusions should be carefully drawn from further studies and future monitoring, Japan's marked health improvement in the past half century may not enjoy an equal parallel in the future.

### Acknowledgements

This study was supported by a Grant-in-Aid for Scientific Research from the Japan Society for the Promotion of Science and a Health and Labour Sciences Research from the Ministry of Health, Labour and Welfare, Japan.

### References

1. Acheson D. Independent Inquiry into Inequalities in Health. London: Stationary Office, 2000.
2. Davey Smith G. Health Inequalities. Bristol: Policy Press, 2003.
3. Drever F, Whitehead M. Health inequalities. London: Stationery Office, 1997.
4. Gordon D, Shaw M, Dorling D, Davey Smith G. Inequalities in health. London: Policy Press, 1999.
5. Wilkinson RG. Income distribution and life expectancy. *BMJ* 1992;304:165-168.
6. Kawachi I, Kennedy BP. The Health of the Nations: Why Inequality Is Harmful to Your Health. New York: The New Press, 2002.
7. Organization WH. World Health Report 2006. Geneva: World Health Organization, 2006.
8. Hasegawa T. Japan: historical and current dimensions of

- health and equity. In: Evans T, Whitehead M, Diderichsen F, Bhuiya A, Wirth M, editors. *Challenging inequalities in health*. New York: Oxford University Press, 2001:90-103.
9. Marmot MG, Davey Smith G. Why are the Japanese living longer? *BMJ* 1989;299:1547-1551.
  10. Takano T, Nakamura K. The national financial adjustment policy and the equalization of health levels among prefectures. *J Epidemiol Community Health* 2001;55:748-754.
  11. Wilkinson RG. *Unhealthy Societies*. New York: Routledge, 1996.
  12. Kariya T. *Education in crisis and stratified Japan*. Tokyo: Yushindo Kobunsha, 2001.
  13. Sato T. *Japan as an inequalities society*. Tokyo: Chuokoron Shinsya, 2000.
  14. Tachibanaki T. *Japanese economy from point of view of household expenditure*. Tokyo: Iwanami Shoten, 2004.
  15. Economic and Social Research Institute. *Annual report on prefectural accounts*. Tokyo: National Printing Bureau, 1956, 1961, 1966, 1971, 1976, 1981, 1986, 1991, 1996, 2000.
  16. Ministry of Health, Labour and Welfare. *Age-adjusted mortality rates by prefecture*. Tokyo: Ministry of Health/Health and Welfare Statistics Association, 1969, 1974, 1978, 1983, 1988, 1992, 1997, 2002.
  17. Ministry of Health, Labour and Welfare. *Prefectural life tables 2000*. Tokyo: Health and Welfare Statistics Association, 2003.
  18. Shigematsu T, Nanjo Z, Yoshinaga K. *Revised edition: Mizushima's life tables by prefecture, Japan, 1921-1961*. Fukuoka: Department of Public Health, Medical School of Fukuoka University, 1996.
  19. Fukuda Y, Nakamura K, Takano T. Municipal socioeconomic status and mortality in Japan: sex and age differences, and trends in 1973-1998. *Soc Sci Med* 2004;59:2435-2445.
  20. Ministry of Health, Labour and Welfare. *Data Base System of Ministry of Health, Labour and Welfare*. Tokyo: Ministry of Health, Labour and Welfare, 2002.
  21. Statistics Bureau, Management and Coordination Agency. *System of Socioeconomic and Demographic Statistics*. Tokyo: Statistical Information Institute for Consulting and Analysis, 2000.
  22. Statistical Information Institute for Consulting and Analysis. *Municipal data of social indicators*. Tokyo: Statistical Information Institute for Consulting and Analysis, 2003.
  23. Wagstaff A, Paci P, van Doorslaer E. On the measurement of inequalities in health. *Soc Sci Med* 1991;33:545-557.
  24. Mackenbach JP, Bos V, Andersen O, Cardano M, Costa G, Harding S, Reid A, Hemström Ö, Valkonen T, Kunst AE. Widening socioeconomic inequalities in mortality in six Western European countries. *Int J Epidemiol* 2003;32:830-837.
  25. Fawcett J, Blakely T, Kunst A. Are mortality differences and trends by education any better or worse in New Zealand? A comparison study with Norway, Denmark and Finland, 1980-1990s. *Eur J Epidemiol* 2005;20:683-691.
  26. Pappas G, Queen S, Hadden W, Fisher G. The increasing disparity in mortality between socioeconomic groups in the United States, 1960 and 1986. *N Engl J Med* 1993;329:103-109.
  27. Ishitani S, Nakamura K, Fukuda Y, Kizuki M, Seino K, Inose T. Critical contribution of sanitation infrastructure and primary medical care to child health status in Japan from 1955 to 2000. *J Med Dent Sci* 2005;52:213-222.
  28. Soukejima S, Kagamimori S. Occupation and life expectancy. *Jpn Med J (Natl Inst Health Jpn)* 2005;4247:129-131.
  29. Ministry of Health, Labour and Welfare. *Life expectancy of Japanese, 2005*. Tokyo: Ministry of Health, Labour and Welfare, 2006.
  30. Forster M, d'Ercole M. *Income distribution and poverty in OECD countries in the second half of the 1990s*. Paris: OECD, 2005.
  31. World Health Organization. *World Health Report, 2000*. Geneva: WHO, 2000.
  32. Matsumoto K, Tanaka M, Maemura T, Hirao T, Hasegawa T, Hasegawa T. Study on supply, demand and maldistribution of obstetricians and gynecologist in Japan. *Hosp Adm (Chic)* 2007;44:93-103.
  33. National Cancer Center. [http://www.ncc.go.jp/jp/cis/admin\\_council/pdf/2nd\\_shiryou-yamasaki.pdf](http://www.ncc.go.jp/jp/cis/admin_council/pdf/2nd_shiryou-yamasaki.pdf)
  34. Imai H, Itou T, Yoshida T, Takuma N, Ishikawa M. Association of numbers of obstetricians, gynecologists and pedestrians with of perinatal indicators. *Jpn Med J* 2005;4246:28-32.
  35. Kawabuchi K. *Apprehension for Japanese health care*. Tokyo: Chikuma Shobo, 2006.
  36. Irohira T, Yamaoka J. *Income inequalities in health care*. Tokyo: Chuo Koron Shinsya, 2005.
  37. Kunst A, Mackenbach J. *Measuring socioeconomic inequalities in health*. Copenhagen: WHO Regional Office for Europe, 1997.
  38. Morris R, Carstairs V. Which deprivation? A comparison of selected deprivation indexes. *J Public Health Med* 1991;13:318-326.
  39. Greenland S, Robins J. Ecological studies, biases, misconceptions, and counter-examples. *Am J Epidemiol* 1994;139:747-760.
  40. Kunst AE. Using geographical data to monitor socioeconomic inequalities in mortality: experiences from Japanese studies. *Int J Epidemiol* 2005;34:110-112.